

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/319016576>

# Release and distress calls of *Rhinella abei* (Baldissera, Caramaschi & Haddad, 2004) and *Rhinella icterica* (Spix, 1824)

Article in *Herpetozoa* · August 2017

CITATIONS

0

READS

90

6 authors, including:



[Vinicius Guerra Batista](#)

Universidade Estadual de Maringá

41 PUBLICATIONS 62 CITATIONS

[SEE PROFILE](#)



[Fabrício Hiroiuki Oda](#)

Universidade Estadual de Maringá

61 PUBLICATIONS 110 CITATIONS

[SEE PROFILE](#)



[Nathane de Queiroz Costa](#)

Amazon Environmental Research Institute (IP...

2 PUBLICATIONS 0 CITATIONS

[SEE PROFILE](#)



[Natan Medeiros Maciel](#)

Universidade Federal de Goiás

39 PUBLICATIONS 290 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Diversity pattern and assemblage structure of anurans from Brazilian Cerrado [View project](#)



IUCN Amphibian Red List Assessments [View project](#)

Release and distress calls of  
*Rhinella abei* (BALDISSERA,  
CARAMASCHI & HADDAD, 2004), and  
*Rhinella icterica* (SPIX, 1824)

Vocalization is the main form of communication in anurans. These animals produce many different types of sounds which have specific meanings in the social context they are used in (WELLS et al. 2007; TOLEDO et al. 2014). The advertisement call is the best known and thus, most frequently described and used call type in taxonomic studies (WELLS et al. 2007; GAMBALE & BASTOS 2014; BATISTA et al. 2015). However, other calls, such as release calls, play important roles in anuran biology. Release calls are emitted by males or females when another frog, either conspecific or not, attempts to mate with them (WELLS et al. 2007; TOLEDO et al. 2014). Like advertisement calls, release calls may vary between species and help distinguish between closely related taxa (BROWN & LITTLEJOHN 1972; GRENAT & MARTINO 2013). Distress call is a defensive scream call emitted by males, females, juveniles, newly metamorphosed as well as larvae of anurans when grasped by potential predators (WELLS 2007;

NATALE et al. 2011; TOLEDO et al. 2014). Loud and explosive notes characterize the distress call, which distinguishes it from other vocalizations (DUELLMAN & TRUEB 1994). Distress calls can be emitted with mouth open or closed (TOLEDO & HADDAD 2009). Calls are stereotyped and also help to identify differences among species (CARVALHO et al. 2013). In the present note, the authors describe the release and distress calls of *Rhinella abei* (BALDISSERA, CARAMASCHI & HADDAD, 2004) (of the *Rhinella crucifer* group) and *Rhinella icterica* (SPIX, 1824) (of the *Rhinella marina* group).

Five male individuals of *R. abei* and three of *R. icterica* were spotted on September 9, 2014, and September 23, 2015. Specimens were found calling from the edge of water bodies, near an Araucaria Forest remnant in the Atlantic Rain Forest domain in the municipality of Campo Largo (25°30' 26.90" S, 49°22' 35.88" W, 905 m above sea level), Paraná state, southern Brazil. The frogs were caught, positioned sitting on the ground, and squeezed in the lateral abdominal region behind the forelimbs, simulating an amplexus. This procedure caused the males to emit release calls. When the males were squeezed the same way lifted off the ground, they emitted distress calls. Recordings were obtained using Marantz® PMD 660 and Tascam® Dr-40 digital recorders connected to a Sennheiser® ME66/K6 directional microphone. Calls were recorded at 44.1 kHz with 16-bit resolution. Bio-acoustic traits were analyzed using Raven Pro 64 1.5 software from the Cornell Lab of Ornithology (BIOACOUSTICS RESEARCH PROGRAM 2014). Spectrograms were produced applying a window size of 256 samples, 75 % overlap, a hop size of 64 samples, DFT of 1024 samples, and Hamming window type. Oscillogram and spectrogram figures were produced using TuneR 1.0 (LIGGES et al. 2013) and Seewave 1.7.3 (SUEUR et al. 2008) packages for R version 3.3.3 (R DEVELOPMENT CORE TEAM 2016). Voucher specimens are housed at the zoological collection of Universidade Federal de Goiás (ZUFG), Goiás state and the herpetological collection of Museu Nacional, Universidade Federal do Rio de Janeiro (MNRJ), Rio de Janeiro state, Brazil (*R. abei*: ZUFG 9884; ZUFG 9885; *R. icterica*: MNRJ 89526; MNRJ

89527). The following acoustic parameters were analyzed: call duration, note duration, note number, internote interval, pulse number, pulse duration, inter-pulse interval, dominant frequency of the call, and lower and upper frequency of the call (see BATISTA et al. 2015; FORTI et al. 2015). Parameters were measured from 87 calls of *R. abei* and 68 calls of *R. ictERICA*. Measurements are presented as the mean (or mode)  $\pm$  standard deviation (minimum – maximum). Call description and terminology follow TOLEDO et al. (2014). Voucher individuals were measured (snout-vent length) using a digital caliper (to the nearest 0.1 mm) and the temperature was taken using a digital thermometer (to the nearest 0.01 °C).

Average snout-vent length (SVL) of *R. abei* males recorded was  $60.01 \pm 2.64$  mm (57–64.02 mm;  $N = 5$ ) and mean air temperature was  $16.7 \pm 0.82$  °C (15.0 – 17.6 °C,  $N = 5$ ). Three types of release call and one of distress call were identified. Release call “A” was emitted by five males of *R. abei* and consists of a single pulsed note (Table 1; Fig. 1A). Release call “B” was emitted by three males and consists of a series of pulsed notes (Table 1; Fig. 1B). In the release call “B” the first ( $0.366 \pm 0.158$  s,  $0.077$ – $0.531$  s,  $N = 10$ ) and last ( $0.355 \pm 0.093$  s,  $0.077$ – $0.208$  s,  $N = 6$ ) notes of the call have longer duration in relation to the middle notes ( $0.058 \pm 0.039$  s,  $0.031$ – $0.187$  s,  $N = 15$ ), which comprise the notes between the first and last notes of the call. Release call “C” was emitted by one male and has a harmonic structure. This call presented three well-defined harmonics (Table 1; Fig. 1C). When emitting release calls, all males of *R. abei* made small body vibrations. Distress calls were emitted by four males of *R. abei* as a loud scream of short duration and harmonic structure (Table 1; Fig. 1D).

*Rhinella ictERICA* specimens recorded had an average SVL of  $96.73 \pm 24.81$  mm (68.09–111.50 mm,  $N = 3$ ) and mean air temperature was  $18.01 \pm 0.70$  °C (17.20–18.41 °C,  $N = 3$ ). The authors identified two types of release call and one of distress call. The release call type “A” was emitted by all males of *R. ictERICA* and consists of a single pulsed note (Table 1; Fig. 2A). The release call type “B” was emitted by two males and

consists of a series of pulsed notes (Table 1; Fig. 2B). While emitting release calls the males of *R. ictERICA* made small body vibrations. The distress call of *R. ictERICA* was emitted by three males and consisted of a single note with a harmonic structure (Table 1; Fig. 2C).

*Rhinella abei* showed a greater release call repertoire than *R. ictERICA*. Such repertoire was composed of single and compound calls. There are no release and distress call descriptions for any other species in the *Rhinella crucifer* group. In the *Rhinella marina* group, the release call is only known for *Rhinella jimi* (STEVAUX, 2002) (GARDA et al. 2010). The release call repertoire of *R. ictERICA* is more extensive than *R. jimi* and the pulse number of the notes is higher. However, the dominant frequency is similar and overlaps in both species. A male of *R. abei* and six males of *R. ictERICA* were evaluated by TOLEDO et al. (2009). However, they did not emit distress calls. In the present work all males that were handled emitted release and/or distress calls. All individuals tested responded to the handling with a series of vibrations, similar with the muscular contractions produced by *R. mirandari-beiroi* (GALLARDO, 1965) (VIEIRA et al. 2014). Vibrations displayed by some anuran species may be important to initiate dismounting of males (WELLS 2007). Such behavior is commonly found among *Rhinella* species (BLAIR 1947; RENGEL 1948; VIEIRA et al. 2014). The release and distress calls recorded by *R. abei* and *R. ictERICA* were emitted with mouths closed as noted in other bufonids (WEBER 1978).

Some of the functions of release calls are to avoid breeding energy expenditure with a misleading amplexus, to warn conspecifics of predator presence and/or to frighten (or surprise) audiotively oriented predators, or even to attract a secondary predator capable of interfering in the predatory process (HÖDL & GOLLMANN 1986; TOLEDO et al. 2014). However, a case of amplexus between a male of *R. ictERICA* and a male of the invasive bullfrog *Lithobates catesbeianus* (SHAW, 1802), indicated that release calls emitted by the amplexed male of *R. ictERICA* were not effective to prevent the interspecific amplexus (THEIS & CALDART 2015). Future researches using experi-




Fig. 1: Release and distress calls of *Rhinella abei* (BALDISSERA, CARAMASCHI & HADDAD, 2004) from Campo Largo, Paraná state, southern Brazil. A, B, C – release call types, D – distress call. Upper graphs: Audiospectrograms. Lower graphs: Oscillograms. Air temperature at time of recording – 17.6 °C. Water temperature – 20.0 °C. Male SVL – 64.2 mm. Unvouchered specimen.




Fig. 2: Release and distress calls of *Rhinella icterica* (SPIX, 1824) from Campo Largo, Paraná state, southern Brazil. A, B – release call types, C – distress call. Upper graphs: Audiospectrograms. Lower graphs: Oscillograms. Air temperature at time of recording – 17.6 °C. Water temperature – 17.2 °C. Male SVL – 110.6 mm. Unvouchered specimen.

Table 1: Acoustic parameters of the release and distress calls of *Rhinella abei* (BALDISSERA, CARAMASCHI & HADDAD, 2004), and *R. ictERICA* (SPIX, 1824). Data are presented as mean  $\pm$  SD (or mode – for parameters presenting only integers (numbers), range and *N* (number of calls analyzed)). *Rhinella abei*: release calls (A, B and C); distress call (D). *Rhinella ictERICA*: release calls (A and B); distress call (C). CD – call duration; NN – note number; ND – note duration; PN – pulse number; PD – pulse duration; Inot – inter-note interval; Ipul – inter-pulse interval; DF – dominant frequency; MaxF – highest frequency of the call; MinF – lowest frequency of the call.

Taxon	Acoustic parameters											
	Call	CD (s)	NN	ND (s)	PN	PD (s)	Inot (s)	Ipul (s)	DF (Hz)	MaxF (Hz)	MinF (Hz)	
<i>Rhinella abei</i>	A	0.442 $\pm$ 0.150 (0.072-0.680) <i>N</i> = 27	–	–	31 (11-82) <i>N</i> = 27	0.007 $\pm$ 0.002 (0.003-0.012) <i>N</i> = 135	–	0.004 $\pm$ 0.004 (0.001-0.040) <i>N</i> = 135	1156.01 $\pm$ 68.08 (1033.6-1205.9) <i>N</i> = 27	1156.01 $\pm$ 68.08 (1033.60-1205.90) <i>N</i> = 27	863.23 $\pm$ 143.57 (516.80-1033.60) <i>N</i> = 27	
	B	1.620 $\pm$ 0.679 (0.585-2.906) <i>N</i> = 10	2 (2-8) <i>N</i> = 10	0.206 $\pm$ 0.366 (0.058-0.161) <i>N</i> = 42	5 (4-71) <i>N</i> = 42	0.007 $\pm$ 0.002 (0.003-0.013) <i>N</i> = 42	0.215 $\pm$ 0.056 (0.142-0.395) <i>N</i> = 32	0.006 $\pm$ 0.004 (0.001-0.021) <i>N</i> = 131	1214.10 $\pm$ 37.12 (125.9-1378.1) <i>N</i> = 42	1533.98 $\pm$ 189.52 (1378.10-1894.90) <i>N</i> = 42	824.40 $\pm$ 775.17 (344.50-1033.60) <i>N</i> = 42	
	C	0.659 $\pm$ 0.037 (0.616-0.682) <i>N</i> = 3	–	–	– (11-21) <i>N</i> = 3	0.008 $\pm$ 0.001 (0.006-0.012) <i>N</i> = 3	–	0.004 $\pm$ 0.002 (0.001-0.007) <i>N</i> = 15	1205.90 (1205.90-1205.90) <i>N</i> = 3	1378.10 (1378.10-1378.10) <i>N</i> = 15	976.17 $\pm$ 99.48 (861.30-1033.60) <i>N</i> = 15	
	D	0.109 $\pm$ 0.198 (0.017-0.615) <i>N</i> = 15	–	–	–	–	–	–	1194.40 $\pm$ 102.25 (1053.60-1378.10) <i>N</i> = 15	1538.90 $\pm$ 322.08 (941.72-143.64) <i>N</i> = 15	941.72 $\pm$ 143.64 (689.10-1205.90) <i>N</i> = 15	
	<i>Rhinella ictERICA</i>	A	0.855 $\pm$ 0.251 (0.477-1.145) <i>N</i> = 9	–	–	33 (17-36) <i>N</i> = 9	0.018 $\pm$ 0.019 (0.009-0.030) <i>N</i> = 145	–	0.011 $\pm$ 0.003 (0.006-0.018) <i>N</i> = 45	650.81 $\pm$ 75.977 (516.80-689.10) <i>N</i> = 9	1684.38 $\pm$ 392.63 (1205.90-2239.50) <i>N</i> = 9	401.97 $\pm$ 172.25 (516.80-172.30) <i>N</i> = 9
		B	0.237 $\pm$ 0.0338 (0.026-1.113) <i>N</i> = 37	5 (3-7) <i>N</i> = 37	0.237 $\pm$ 0.338 (0.026-1.113) <i>N</i> = 37	1 (1-33) <i>N</i> = 37	0.022 $\pm$ 0.012 (0.010-0.044) <i>N</i> = 35	0.284 $\pm$ 0.057 (0.117-0.389) <i>N</i> = 29	0.014 $\pm$ 0.004 (0.005-0.019) <i>N</i> = 35	786.87 $\pm$ 105.20 (689.10-1205.90) <i>N</i> = 37	1569.02 $\pm$ 324.29 (1205.90-2239.50) <i>N</i> = 37	428.34 $\pm$ 138. (172.30-516.80) <i>N</i> = 37
		C	0.061 $\pm$ 0.031 (0.029-0.122) <i>N</i> = 18	–	–	1 (1-2) <i>N</i> = 18	0.045 $\pm$ 0.025 (0.016-0.075) <i>N</i> = 8	–	0.008 $\pm$ 0.003 (0.005-0.011) <i>N</i> = 4	746.51 $\pm$ 132.11 (689.10-1205.90) <i>N</i> = 18	1512.10 $\pm$ 274.34 (861.30-2067.20) <i>N</i> = 18	507.23 $\pm$ 124.96 (172.30-689.10) <i>N</i> = 18

ments could be conducted in order to better understand the role of release and distress calls. The taxonomic status of some bufonid species is problematic due their similar external morphology and on account of hybridization zones between sympatric species such as e.g., *Rhinella ornata* (SPIX, 1824), and *R. abei* (THOMÉ et al. 2012). Basic natural history information is fundamental to evaluate the process shaping the ecology of the species and to differentiate the species-specific characters of important taxonomic value.

**ACKNOWLEDGMENTS:** The authors thank the Postgraduate Program on Ecology of Continental Aquatic Environments (PROEX/PEA) for the logistical support. Thanks to Priscila C. SILVEIRA for critical reading and valuable comments on the manuscript. V. G. BATISTA and F. H. ODA would like to thank The Coordination for the Improvement of Higher Education Personnel (CAPES) by fellowships.

**REFERENCES:** BATISTA, V. G. & GAMBALE, P. G. & LOURENÇO-DE-MORAES, R. & CAMPOS R. M. & BASTOS, R.P. (2015): Vocalizations of two species of the *Hypsiboas pulchellus* group (Anura: Hylidae) with comments on this species group.- North-Western Journal of Zoology, Oradea; 11: 253-261. BIOACOUSTICS RESEARCH PROGRAM (2014): Raven Pro: Interactive sound analysis software. Version 1.5. The Cornell lab of ornithology, Ithaca, New York. WWW resource, information and software available at < <http://www.birds.cornell.edu/raven> >. BLAIR, A. P. (1947): The male warning vibration in *Bufo*.- American Museum Novitates, New York; 1344: 1-7. BROWN, L. E. & LITTLEJOHN, M. J. (1972): Male release call in the *Bufo americanus* group; pp. 310-323. In: BLAIR, W. F. (Ed.): Evolution in the genus *Bufo*. Austin (University of Texas Press). CARVALHO, T. R. & GIARETTA, A. A. (2013): Taxonomic circumscription of *Adenomera martinezi* (BOKERMANN, 1956) (Anura: Leptodactylidae: Leptodactylinae) with the recognition of a new cryptic taxon through a bioacoustic approach.- Zootaxa, Auckland; 3701: 207-237. DUELLMAN, W. E. & TRUEB, L. (1994): Biology of amphibians. Baltimore and London (Johns Hopkins University Press), pp. 670. FORTI, L. R. & MÁRQUEZ, R. & BERTOLUCI, J. (2015): Advertisement call of *Dendropsophus microps* (Anura: Hylidae) from two populations from southeastern Brazil.- Zoologia, Curitiba; 32: 187-194. GAMBALE, P. G. & BASTOS, R. P. (2014): Vocal repertoire and bioacoustic analyses in *Physalaemus cuvieri* (Anura, Leptodactylidae) in southern Brazil.- Herpetological Journal, London; 24: 31-39. GARDA, A. A. & SÃO PEDRO, V. A. & LION, M. B. (2010): The advertisement and release calls of *Rhinella jimi* (Anura, Bufonidae).- South American Journal of Herpetology, São Paulo; 5: 151-156. GRENAT, P. R. & MARTINO, A. L. (2013): The release call as a diagnostic character between cryptic related species *Odontophrynus cordobae* and *O. americanus* (Anura: Cycloramphidae).- Zootaxa, Auckland; 3635: 583-586. HÖDL, W. & GOLLMANN, G. (1986): Distress calls in Neotropical frogs.- Amphibia-Reptilia, Leiden; 7: 11-21. LIGGES, U. &

KREY, S. & MERSMANN, O. & SCHNACKENBERG, S. (2013): Tuner: Analysis of music. WWW resource, information and software available at < <http://r-forge.r-project.org/projects/tuner/> >. [last accessed on January 10, 2015]. NATALE, G. S. & ALCALDE, L. & HERRERA, R. & CAJADE, R. & SCHAEFER, E. F. & MARANGONI, F. & TRUDEAU, V. L. (2011): Underwater acoustic communication in the macrophagic carnivorous larvae of *Ceratophrys ornata* (Anura: Ceratophryidae).- Acta Zoologica : Royal Swedish Academy of Sciences etc.; Oxford; 92: 46-53. R DEVELOPMENT CORE TEAM (2016): R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0. WWW resource, information and software available at < <http://www.R-project.org/> >. RENGEL, D. (1948): Sobre la vibración preventiva ("warning vibration") en los sapos machos del norte argentino.- Acta Zoológica Lilloana, Tucumán; 6: 279-282. SUEUR, J. & AUBIN, T. & SIMONIS, C. (2008): Seewave: a free modular tool for sound analysis and synthesis.- Bioacoustics, Abingdon; 18: 213-226. THEIS, T. F. & CALDART, V. M. (2015): Multiple interspecific amplexus between a male of the invasive Bullfrog *Lithobates catesbeianus* (Ranidae) and two males of the Cururu toad *Rhinella icterica* (Bufonidae).- Herpetology Notes, Braunschweig; 8: 448-451. THOMÉ, M. T. C. & ZAMUDIO, K. R. & ALEXANDRINO, J. (2012): Delimiting genetic units in Neotropical toads under incomplete lineage sorting and hybridization.- BMC Evolutionary Biology, London etc.; 12: 242. TOLEDO L. F. & C. F. B. HADDAD (2009): Defensive vocalizations of Neotropical anurans.- South American Journal of Herpetology, São Paulo; 4: 25-42. TOLEDO, L. F. & MARTINS, I. A. & BRUSCHI, D. P. & PASSOS, M. A. & ALEXANDRE, C. & HADDAD, C.F.B. (2014): The anuran calling repertoire in the light of social context.- Acta Ethologica, Berlin, Heidelberg; 18: 87-99. VIEIRA, R. R. S. & BASTOS, R. P. & GAMBALE, P. G. (2014): The release call of *Rhinella mirandaribeiroi* (GALLARDO, 1965) (Anura: Bufonidae).- Herpetology Notes, Braunschweig; 7: 543-545. WEBER, E. (1978): Distress calls of *Bufo calamita* and *B. viridis* (Amphibia: Anura).- Copeia, Washington; 1978: 354-356. WELLS, K. D. (1977): The social behavior of anuran amphibians.- Animal Behavior, Amsterdam; 25: 666-693.

**KEY WORDS:** Amphibia: Anura: Bufonidae: *Rhinella abei*, *Rhinella icterica*, *Rhinella crucifer* group, *Rhinella marina* group; acoustic behavior, vocalizations, acoustic parameters, call repertoire, Atlantic Rain Forest domain, Brazil

SUBMITTED: April 12, 2016

**AUTHORS:** Vinicius G. BATISTA (Corresponding author < [vinicius.guerrabatista@gmail.com](mailto:vinicius.guerrabatista@gmail.com) >)<sup>1</sup>, Fabrício H. ODA<sup>2</sup>), Diogo F. DO AMARAL<sup>3</sup>), Nathane DE Q. COSTA<sup>4</sup>), Natan M. MACIEL<sup>4</sup>) & Rogério P. BASTOS<sup>4</sup>)

<sup>1</sup>) Universidade Estadual de Maringá (UEM), Programa de Pós-graduação em Ecologia de Ambientes Aquáticos Continentais. Av. Colombo, 5790, CEP 87020-900, Maringá, PR, Brazil.

<sup>2</sup>) Centro Universitário Cesumar (UniCesumar), Programa de Pós-Graduação em Promoção da Saúde. Avenida Guedner 1610, Jardim Aclimação, CEP 87050-390, Maringá, PR, Brazil.

<sup>3</sup>) Programa de Pós-Graduação em Conservação de Recursos Naturais do Cerrado. Rodovia Geraldo

---

Silva Nascimento, Km 2,5—Zona Rural, CEP 75.790-000, Urutai, GO, Brazil.

<sup>4)</sup> Universidade Federal de Goiás (UFG), Instituto de Ciências Biológicas, Departamento de Ecologia, Laboratório de Herpetologia e Comportamento Animal, Campus Samambaia, 74001-970, Cx. Postal 131, Goiânia, GO, Brazil.